



# ENDO GARD



## ENDO GARD

### Palatable Allwormer for Dogs

- ▶ **Kills and controls ALL intestinal worms**  
**Endogard** is for the treatment and prevention of roundworm, hookworm, whipworm and tapeworm, including hydatids. Endogard is safe to use with any heartwormer for dogs on the market.\* It is an integral part of an effective parasite control program for dogs.
- ▶ **Tasty Liver Flavour**  
Dogs will love the liver flavoured chewable tablet so administration is easy.
- ▶ **Flexible Dosing**  
Endogard chewable tablets are quarter scored to allow for flexible dosing in multi-pet households.

## INTESTINAL WORMER

### PROPERTIES

Broad Spectrum  
Tasty Liver flavour  
Easy to administer  
Safe

### COMPOSITION

Contains praziquantel (5mg/kg) and oxbendazole (22.5mg/kg).

### PRESENTATION

Small dog (0-5kg) tablet  
(25mg praziquantel and 112.5mg oxbendazole)

Medium dog (5-10kg) tablet  
(50mg praziquantel and 225mg oxbendazole)

Large dog (10-20kg) tablet  
(100mg praziquantel and 550mg oxbendazole)

Available in cartons and bulk packs

### INDICATIONS

Treatment and control of roundworm (*Toxascaris leonina*, *Toxacara canis*), hookworm (*Ancylostoma caninum*, *Ucinaria stenocephala*), whipworm (*Trichuris spp*) and tapeworm (*Dipylidium caninum*, *Taenia spp*), including hydatids (*Echinococcus granulosus*) in dogs. Does not control heartworm.

### DIRECTIONS FOR USE

**Young puppies:** Treat according to weight at 2, 4, 8 and 12 weeks of age.

**Adult Dogs:** Treat according to weight every three months. If your pet is a breeding bitch, or located in a known hydatid area or if whipworm is a problem, more frequent administration is required.

# ENDOARD



## SAFETY OF ENDOARD WITH COMMONLY USED INSECTICIDES

Monash University conducted a confirmatory safety study with **Virbac Endogard Palatable Allwormer** when administered concurrently with insecticides\*.

### Method

- ▶ A total of 16 pups were used (9 greyhounds and 7 crossbred pups) ranging in age from 3-7 months.
- ▶ These pups were divided into 3 groups, 2 of which were given a combination of commonly used insecticides as outlined below, as well as the test material.
- ▶ The third group was only given **Virbac Endogard Palatable Allwormer**.
- ▶ The group of actives in group 2 covered the routine chemicals used as dog ectoparasiticides.
- ▶ The insecticides used were applied as per manufacturer's recommendations.

| Timetable for insecticide and Endogard administration |  |          |  |             |
|---|--|----------|--|-------------|
|   | Day 0  | Day 14   | Day 28   | Day 35      |
| Group 1   | HeartGuard®<br>Program®<br>Frontline® &<br><b>Endogard</b> | Endogard | Program®<br>Frontline®<br>Proban® &<br><b>Endogard</b> | Final Check |
| Group 2   | Dimmitrol®<br>Preventic®<br>Care® &<br><b>Endogard</b>     | Endogard | Care® &<br><b>Endogard</b>                             | Final Check |
| Group 3   | Endogard   | Endogard | Endogard   | Final Check |

- ▶ All dogs were observed daily during the trial.
- ▶ They were also individually examined at 0,6,24 and 48 hours after each administration.

### Results

- ▶ No abnormal developments were noted at any time point, examination or observation.

### Conclusion

This study confirms that there was no evidence of deleterious effects on the pups. Virbac Endogard Palatable Allwormer is safe to use in dogs under most conditions, particularly as the trial conditions would normally exceed routine use of insecticides, heartworm treatment and gastrointestinal worming treatment.

\* Data on File, Study No.: 97/15MM/KB (Melbourne, Monash University Animal Services)