

# Stampede®

## Stampede Pour-On Lousicide for Cattle and Sheep.

Stampede is a pour-on formulation designed for both cattle and sheep, containing the revolutionary active constituent diflubenzuron, to provide long lasting control against susceptible biting and sucking lice from a single treatment in cattle, and body lice in sheep.

### WHAT IS STAMPEDE?

- Stampede is the only cattle and sheep lousicide to contain Insect Growth Regulator (IGR) technology.
- Stampede contains 20g/L diflubenzuron, an IGR that prevents nymph stages of lice from developing, due to its action against the enzyme chitin synthetase.
- Stampede stays active on the animals for months after treatment reducing the likelihood of retreatment during the season.

### WHY CHOOSE STAMPEDE?


- ✓ **Cattle lice** - highly effective and provides long term control against biting and sucking lice in cattle.
- ✓ **Sheep lice** - highly effective and provides long term control against body lice in sheep.
- ✓ **Fast acting.**
- ✓ **IGR** - Insect Growth Regulator technology.
- ✓ **Safe and easy to use.**
- ✓ **Nil milk** withhold for cattle.
- ✓ **Short WHP and ESI** - for both cattle and sheep.
- ✓ **Rainfast.**
- ✓ **Versatile** - don't double up on buying separate products for cattle and sheep.
- ✓ **Solvent based.**
- ✓ **Economical** - dose rates for cattle and sheep allow both to be economically treated.



### WHEN TO USE STAMPEDE

- Treat cattle for lice after the first cold snap in autumn, when eggs are likely to be hatching.
- Consider treating bulls prior to joining with an additional treatment if necessary.
- Treat sheep up to 7 days off shears.
- Treat unshorn lambs at foot up to 3 months old and greater than 10 kg.
- Ewes due to lamb post shearing should be shorn and treated at least 4 weeks prior to lambing.
- Don't mix treated animals with untreated (clean) animals for at least 6-8 weeks.

Typical seasonal profile for lice in cattle (Schematic diagram)



**Prevention is better than cure.**  
Treat cattle with Stampede before lice numbers build up.

# Stampede®

## QUICK REFERENCE GUIDE

### INDICATIONS:

Treatment of body lice (*Bovicola bovis*) in sheep and biting lice (*Bovicola bovis*) and sucking lice (*Linognathus vituli*, *Haematopinus eurysternus* and *Solenopotes capillatus*) in cattle.

### RESTRAINTS:

- Do not re-treat cattle within 21 days.
- Do not apply to wet sheep.
- Avoid use in extreme heat (>40 deg C).
- Do not use on lactating or pregnant ewes where milk or milk products may be used for human consumption.
- Do not use on sheep that have not been cleanly shorn.

### CONTRAINDICATIONS:

Do not administer orally, intravenously or intramuscularly.

### DOSAGE & ADMINISTRATION:

For topical use only. A representative sample of animals should be weighed before treatment either with scales or a weigh band. Dose according to the heaviest animal by live weight in the group. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

#### CATTLE

Liveweight (kg)	Dose Volume (mL)	5L treats	20L treats
Under 100	25	200	800
101-200	40	125	500
201-300	50	100	400
301-400	60	83	333
401-500	70	71	285
Over 500	Add 10mL/100kg		

#### SHEEP

Liveweight (kg)	Dose Volume (mL)	5L treats	20L treats
10-20	12	416	1666
21-30	15	333	1333
31-55	20	250	1000
56-75	25	200	800
Over 75	Add 5mL/10kg		

Treat all animals in the mob. Check the accuracy of the applicator before and during treatment.


### APPLICATION:

#### CATTLE

**Use only with the Stampede applicator for cattle.** Apply along the midline of the back between the wither and the base of the tail.



For better coverage of larger cattle (over 500 kg), the calculated dose may be applied in an "I" shape with stripes across the shoulders and across the base of the tail, joined by a stripe along the midline of the back between the wither and the base of the tail.



#### SHEEP

**Use only with the Stampede T-bar applicator for sheep.** Apply along the midline of the back from the poll to the base of the tail. Do not deviate from the spine.



### CLEAN UP:

Applicators must not be washed out with water. Pump the applicator dry, then lubricate with vegetable oil. Dismantle applicator and wipe with clean cloth.

### PACK SIZE PRESENTATIONS:

- 5 L backpack and 20 L drum

### COMPOSITION:

- 20 g/L Diflubenzuron in 300 g/L N-methyl-2 pyrrolidone

### STORAGE:

Store below 30 degrees Celsius (room temperature) in original container. Do not store for prolonged periods in direct sunlight.

### WITHHOLDING PERIODS:

- Meat: Cattle WHP: 10 days  
Cattle ESI: 10 days  
Sheep WHP: 7 days  
Sheep ESI: 7 days
- Milk: Cattle: Nil  
Sheep: Do not use in sheep producing milk for human consumption
- Wool: Sheep: 6 months

### APVMA No:

- 49806

Stampede®