

Olive Spray Timing Chart For SACOA BIOPEST® Paraffinic Oil

This chart provides a guide for the timing of sprays for pests and diseases that can be managed with BIOPEST®. Refer to How To Use section on website (www.sacoa.com.au/olives.htm) or in the BIOPEST® Olive booklet for details on rates etc (contact SACOA for a copy - 08 9386 7666).

Note: This is a guide only. Status of each pest varies regionally and monitoring is critical to avoid unnecessary or poorly timed sprays.

PESTS	MONTH											
	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun
SCALE												
Soft Brown/Black Scale.	M	M	M	D	D	D	M	D	D	M	M	M
Olive scale.	M	M	M	D	D	D	M	D	D	D	D	M
California Red Scale.	M	M	M	M	M	D	D	D	D	D	M	M
Oleander scale.	M	M	M	M	D	D	D	D	D	D	M	M
BUGS												
Olive Lace Bug.	M	M	D	D	D	D	D	D	D	D	M	M
MITES												
Mites.	-	-	M	D	D	D	M	M	D	D	D	-
DISEASES												
Peacock Spot	-	D	D	M	M	-	-	M	D	D	D	M
OTHER												
Snails/Slugs.	-	-	D	D	M	-	-	M	D	D	-	-
Ants.	M	M	M	D	D	D	D	D	D	D	M	D

M - Pest Monitoring Period

D - Pest Activity/Damage Period

- Best Time For Spraying

Timings Will Vary

This chart provides a suggested approach to IPM program timings.

The timings of insect activity noted here are a guide only, as activity can vary with different seasons.

In general activity commences earlier the further north the planting. Hence regular monitoring of groves is essential.